

Entropy Analysis of Stochastic Processes at Finite Resolution

D.M. Tavares and L.S. Lucena

*International Center for Complex Systems and
Departamento de Física Teórica e Experimental - UFRN
Natal-RN 59078-970 Brazil*

The time evolution of complex systems usually can be described through stochastic processes. These processes are measured at finite resolution, what necessarily reduces them to finite sequences of real numbers. In order to relate these data sets to realizations of the original stochastic processes (to any functions, indeed) it is obligatory to choose an interpolation space (for example, the space of band-limited functions). Clearly, this choice is crucial if the intent is to approximate optimally the original processes inside the interval of measurement. Here we argue that discrete wavelets are suitable to this end. The wavelet approximations of stochastic processes allow us to define an entropy measure for the order-disorder balance of evolution regimes of complex systems, where order is understood as confinement of energy in simple local modes. We calculate exact results for the fractional Brownian motion (fBm), with application to Kolmogorov K41 theory for fully developed turbulence.

PACS numbers: 05.45.Tp, 05.65.+b, 89.75.Fb, 02.50.-r

Many physical systems investigated at present have a complex evolution in time. Frequently, the major information we can obtain on their dynamics comes from time series of noisy appearance [1, 2, 3, 4, 5]. These series are samples at finite resolution of subjacent stochastic processes, whose properties are better investigated through a multi-resolution approach, because the realizations of the mentioned stochastic processes are singular everywhere functions. This is, for example, the case of $1/f^\alpha$ noises from self-organized systems [6, 7, 8, 9, 10]. Thus, singularities are common, and should be interpreted as details that influence function variation at all scales [11, 12]. In this paper we propose a method for entropy analysis of arbitrary complex time series that takes advantage of this theoretical standpoint. Moreover, we account for the fact that measurements are made at finite resolution, considering the consequences of sampling, what is not fully accomplished in previous approaches. In general, other formulations are strongly influenced by information-theoretical arguments, applied to the analysis of chaotic behavior [13]. We see difficulties in two main aspects. First, the notion of complexity that is employed in these formulations is based on entropy rates, like the Kolmogorov-Sinai entropy, that measure the degree in which information on initial conditions is lost when the systems evolve. This gives a scale of complexity ranging from zero (non-chaotic deterministic case) to infinite (stochastic case). In this scale, finite values quantify the complexity of deterministic chaos. As a consequence, the problem of defining a proper complexity estimator for stochastic processes is substituted by the statistical investigation of chaotic deterministic dynamical systems [14]. Second, these formulations do not consider the relationship that scaling has with disorder [15]. Our approach is tailored to face these difficulties. Here, we begin by assuming that the stochastic processes are supported on

the real axis, and measured at a discrete set of points with a sampling interval τ , during a time T . The resolution of the measurement is $N = T/\tau$, and for convenience we make $T = 1$, and $\tau = 1/N = 2^{-J}$ (we will work in these units). Such measurement results in a loss of information, which depends on two factors: The resolution, and the interpolation space in which the stochastic processes will be projected [16]. The idea is that the mere sampled values say almost nothing about a function. Much more information is conveyed through the hypothesis on how the function varies between the sampled values. This regularity hypothesis, usually implicit when we draw smooth curves between data points, is an essential element of the theory. Without this hypothesis, no information found on the discrete and finite data sets can be attributed to the subjacent model, that is assumed to hold on the continuous support. Now, in the time-scale (time-frequency) domain, it makes sense to search a representation *i*) that is minimally affected by the created end singularities, *ii*) that provides an interpolation based on multi-scale approximation of the actual singularities, and *iii*) that deals with the resolution N as a direct experimental parameter. The first and second requirements are the most crucial, and establish the way in which the energy (L^2 -norm squared) is assumed to be distributed inside the interval of measurement, so that this distribution corresponds to the singularity structure of the process, seen at a finite resolution. As a supplementary condition, the best is that the algorithmic complexity grows only linearly with the length of the time series. These requirements are met if we project the stochastic processes in a discrete wavelet space.

Wavelets are associated with time-scale representations. Let $|F\rangle$ be a vector in a Hilbert space. There are bases $\{|\psi_{jn}\rangle\}$ such that $|F\rangle$ can be expanded as




FIG. 1: The Haar wavelet decomposition of a stochastic process sampled at the finite resolution $N = 2^{15}$. The details are presented from scale the 2^{-14} to the scale 2^{-6} . Note how the energy accumulates at the small scale details. This causes the high entropy $S = 12.52$ as compared to the maximum $S_{\max} = 15$.

$$|F\rangle = \sum_{j=-\infty}^{j=\infty} \sum_{n=-\infty}^{n=\infty} \langle \psi_{jn} | F \rangle |\psi_{jn}\rangle, \quad (1)$$

the form of the basis functions being $\psi_{jn}(t) = \frac{1}{\sqrt{2^j}} \psi\left(\frac{t-2^j n}{2^j}\right)$. These functions are called discrete wavelets, and are square integrable with zero mean, and may have p vanishing moments. The discrete wavelet transform of F to this basis is written $D_{jn}F \equiv \langle \psi_{jn} | F \rangle$, and gives information on the behavior of the function F at scale 2^j and time $2^j n$ [12].

Confining the function in expression (1) to a finite observation window, and measuring at finite resolution, we obtain the approximation

$$|f_J\rangle = \langle \phi | f \rangle |\phi\rangle + \sum_{j=-J+1}^{j=0} \sum_{n=0}^{n=2^{-j}-1} \langle \psi_{jn} | f \rangle |\psi_{jn}\rangle, \quad (2)$$

where f is the restriction of F to $[0, 1]$. The function ϕ is called the scaling function, having the properties of a smoothing kernel [12]. We are interested in the fluctuations, so, without any loss, we employ in our analysis the transform $f \rightarrow f - \langle f \rangle$, getting from f a version with null time-average. At this point we are ready to express an energy and an entropy for the time series, that are also resolution dependent approximations of the energy density and entropy density for the ideal stochastic process. The energy of f_J is defined by the ensemble average of the L^2 -norm squared,

$$E(J) \equiv \overline{\langle f_J | f_J \rangle}, \quad (3)$$

that is directly determined with the simple formula $E(J) = \overline{\sum_{i=1}^{i=N} f_i^2}$, where f_i are the sampled values of

f . The entropy has the definition

$$S(J) \equiv - \sum_{j=-J+1}^{j=0} \sum_{n=0}^{2^{-j}-1} P_{jn} \log_2 P_{jn} - P_\phi \log_2 P_\phi, \quad (4)$$

with $P_{jn} \equiv \overline{|D_{jn}F|^2}/E(J)$, and $P_\phi \equiv \overline{|\langle \phi | f \rangle|^2}/E(J)$. This expression has similarities with the basis entropy cost, using the language of reference [12]. In the limit of a single pulse localized around t and with duration $\delta t \sim k\tau$, with k a natural number, only very few wavelet coefficients around the scale $k\tau$ and localized near t will be appreciably different from zero. This characterizes low entropy processes. On the other hand, the white noises (WN) in which the energy is, in average, equally distributed on all wavelets (the same power dissipation at all times and scales) have the maximum entropy $S_{\max} = J$. This entropy functional is useful to quantify the balance between order and disorder (complexity) for different regimes of evolution of complex systems. Actually, we expect that the level of disorder in a process mainly depends on the energy distributed into the details. Consider, for example, the interaction between two surfaces that slip one over the other. When there is no friction we expect no sound. But if we introduce roughness, the energy will be dissipated in microscopic collisions, originating a sound that is classified as a noise, due to the excess of details. Indeed, the intervals between such collisions are so small that no measurement will capture them. The process will continue a noise for ‘high’ resolution instruments, like it is for our ‘low’ resolution ears. In FIG. 1 we see the wavelet decomposition of the sound caused by a piece of plastic pressed and pushed on a rough table. The calculated entropy was $S = 12.52$ for a maximum of $S_{\max} = 15$. The random small scale details have amplitudes comparable to that of the function during the whole observation, constituting important components. The great amount of energy distributed on the multitude of details causes this high entropy. This is why we suggest that the proposed entropy is physically appropriate.

A paradigmatic model for the self-affine processes is the fractional Brownian motion (fBm), introduced by Mandelbrot and Van Ness in the realm of stochastic processes with $1/f^\alpha$ spectra [17]. The fBm is a mon-parametric stochastic process B_H with null ensemble average, Gaussian increments, and whose correlation function has the general form:

$$\overline{B_H(t) B_H(s)} = \frac{\sigma_H^2}{2} \left(|t|^{2H} + |s|^{2H} - |t-s|^{2H} \right), \quad (5)$$

with $\sigma_H^2 \equiv \overline{B_H^2(1)}$. The parameter of the process is the Hurst exponent $H \in (0, 1)$, which controls the nature of the correlations. The variance of B_H depends on time according to $\overline{B_H^2} = \sigma_H^2 |t|^{2H}$, that reduces to the Brownian motion form when $H = 1/2$, representing the uncorrelated case. In this process, for $H < 1/2$ the increments




FIG. 2: The solid line is the theoretical fBm entropy at resolution $N = 2048$. We see that the SRA algorithm gives a very good approximation to the theoretical prediction. Two lines represent the effect of Gaussian white noise superposition. The (visually) straight dotted line at the upper part gives the entropy of simulated pure Gaussian white noise.

present infinite range anti-correlations, while for $H > 1/2$ there are infinite range correlations. The Fourier spectrum of fBm follows the power law

$$S(f) \sim \frac{1}{f^{2H+1}}. \quad (6)$$

where $C(H, J) = [1 - 2^{-2HJ}] / (1 - 2^{-2H})$.

This entropy density is universal and exact for the fBm processes sampled without rounding error to a given finite resolution. It is very useful since for a great number of practical applications there are just low resolution data, what makes mandatory that the theoretical predictions take resolution into account explicitly. In the limit $J \rightarrow \infty$ the entropy diverges at $H = 0$ as $S(H, \infty) \sim 1/2H \ln 2$. For other values of H the entropy is finite at infinite resolution. At finite resolution $S(0, J) = (J - 1)/2 + \log_2(J)$.

The FIG. 2 shows a plot of fBm theoretical entropy for the resolution $N = 2048$. The theoretical result is compared with $S(H, J)$ calculated from samples simulated with the Successive Random Additions (SRA) algorithm [18]. It shows also how Gaussian white noise (WN) affects the entropy, clarifying what is to be expected from noisy data. For comparison, the dots at the upper part of the figure show the entropy for the simulated pure Gaussian white noise, whose average value is $S = 10.98 \sim 11$.

In the Haar wavelet basis one finds exact and simple expressions for the resolution dependent energy and entropy densities of fBm. The Haar wavelets are periods of square waves written as

$$h_{jn}(t) = \begin{cases} \frac{1}{\sqrt{2^j}}, & 2^j n \leq t < 2^j (n + 1/2) \\ -\frac{1}{\sqrt{2^j}}, & 2^j (n + 1/2) \leq t < 2^j (n + 1) \end{cases} \quad (7)$$

For Haar wavelets one proves, using the expression (5), that

$$\overline{|\Delta_{jn} b_{H,J}|^2} = \frac{\sigma_H^2 (1 - 2^{-2H})}{(2H + 1)(2H + 2)} 2^{(2H+1)j}, \quad (8)$$

where $\Delta_{jn} f \equiv \langle h_{jn} | f \rangle$, and b_H indicates restriction of B_H to $[0, 1]$. This expression gives the scale dependent power spectrum in the Haar basis, from which one can calculate the energy density of B_H

$$E(H, J) = \overline{\langle b_{H,J} | b_{H,J} \rangle} = \frac{\sigma_H^2 (1 - 2^{-2HJ})}{(2H + 1)(2H + 2)}, \quad (9)$$

depending on σ_H , that must be determined experimentally. The entropy density has the final form

$$S(H, J) = 2^{-2H} (2H + 1) \frac{1 - 2^{-2H(J-1)} J + 2^{-2HJ} (J - 1)}{(1 - 2^{-2H}) [1 - 2^{-2HJ}]} + \log_2 C(H, J), \quad (10)$$

In the dotted line we see the effect of a superposed noise with 10% of the amplitude of the fBm. The dash-dotted line shows this effect when the percentage grows to 50%. The entropies for all simulated processes were calculated from 100 samples. The drastic diminution of entropy when H increases happens because the details (roughness) become weaker. There is very good agreement between theoretical prediction and simulation.

In conclusion, we propose that the entropy measure (4) is suitable to estimate the complexity of time series by revealing the degree in which *detail modes* are excited in a process. These detail modes are easily represented as localized oscillations or wavelets, which in turn are very effective mathematical instruments for time-scale analysis. Considering such phenomena as fluid or scalar turbulence [19, 20], that are characterized by fluctuating cascades from high inertial scales to low dissipative scales, or plasma turbulence [21], in which some scales are most important for the energy transfer, the proposed entropy seems a natural measure of the intrinsic

order-disorder balance. In such phenomena, many structures (vortices, convective cells) appear at intermediate scales, that should be detected as excitations of intermediate scale modes, lowering the entropy, as compared with noises that are completely determined by microscopic dissipation (whose characteristics are indicated in FIG. 1). For example, if a theory with the same correlations of Kolmogorov K41 theory [22] is valid, implying $H = 1/3$, the time series of velocity increments would lead to the entropy $S(1/3, \infty) = 4.27\dots$, a number significantly small, compared to the case in FIG. 1. What we learn from the fBm analysis presented here suggests that the measurement of the entropy is a strong method of characterization. Furthermore, it is interesting to obtain experimental curves of entropy for stochastic processes depending on parameters, or in situations where the entropy may change with time. This method has given results (which will be published soon) in three situations: In the analysis of the sound of lungs, in the analysis of heart beat series, and in the analysis of global positioning system errors caused by equatorial spread F in ionospheric plasma. We observe that such entropy measurements could give important information on the non-equilibrium approach to self-organized regimes [23], as well as, on self-organized regimes made unstable, for example, by diffusion [24].

We acknowledge the CAPES, CNPq, FINEP and CT-PETRO for financial support.

-
- [1] G. Consolini, T. Chang, *Complexity, magnetic field topology, criticality, and metastability in magnetotail dynamics*: Journal of Atmospheric and Solar-Terrestrial Physics **64**, 541-549 (2002).
 - [2] P.A. Politzer, *Observation of Avalanchelike Phenomena in a Magnetically Confined Plasma*: Phys. Rev. Lett. **84**, 1192-1195 (2000).
 - [3] L. Santi, J.C. Denardin, M.R. Dotto, L.F. Schelp, and R. Sommer, *Barkhausen noise measurements in materials with vanishing magnetoelastic anisotropies*: J. Applied Phys. **91**, 8201-8203 (2002).
 - [4] K.R. Sreenivasan, *On Local Isotropy of Passive Scalars in Turbulent Shear Flows*: Proc. Roy. Soc. London A, **434**, Issue 1890, 165-182 (1991).
 - [5] R. Yulrnyetev, P. Hänggi, and F. Gafarov *Quantification of heart rate variability by discrete nonstationary non-Markov stochastic processes*: Phys. Rev. E **65**, 046107 (2002).
 - [6] P. Bak, C. Tang, K. Wiesenfeld, *Self-Organized Criticality: An explanation of 1/f Noise*: Phys. Rev. Lett. **59**, 381-384 (1987).
 - [7] G. Durin, and S. Zapperi, *Scaling Exponents for Barkhausen Avalanches in Polycrystalline and Amorphous Ferromagnets*: Phys. Rev. Lett. **84**, 4705-4708 (2002).
 - [8] J.S. Urbach, R.C. Madison, and J.T. Markert, *Interface Depinning, Self-Organized Criticality, and the Barkhausen Effect*: Phys. Rev. Lett. **75**, 276-279 (1995).
 - [9] P.J. Cote, and L.V. Meisel, *Self-Organized Criticality and the Barkhausen Effect*: Phys. Rev. Lett. **67**, 1334-1337 (1991).
 - [10] B.A. Carreras, B. van Milligen, M.A. Pedrosa, R. Balbín, C. Hidalgo, D.E. Newman, E. Sánchez, M. Frances, I. García-Cortés, J. Bleuel, M. Endler, S. Davies, and G.F. Matthews, *Long-Range Time Correlations in Plasma Edge Turbulence*: Phys. Rev. Lett. **80**, 4438-4441 (1998).
 - [11] J.F. Muzy, E. Bacry, and A. Arnéodo, *The multifractal formalism revisited with wavelets*: Int. J. of Bifurcation and Chaos **4**, 245-302 (1994).
 - [12] S. Mallat, *A Wavelet Tour of Signal Processing* 2nd. ed. (Academic Press, Boston, 1998).
 - [13] C. Bandt and B. Pompe, *Permutation Entropy: A Natural Complexity Measure for Time Series*: Phys. Rev. Lett. **88**, 174102 (2002).
 - [14] J.P. Eckmann and D. Ruelle, *Ergodic theory of chaos and strange attractors*: Rev. Mod. Phys. **57**, 617-656 (1985).
 - [15] M. Costa, A.L. Goldberger, and C.K. Peng, *Multiscale Entropy Analysis of Complex Physiologic Time Series*: Phys. Rev. Lett. **89**, 068102-1 (2002).
 - [16] M. Unser, *Sampling - 50 Years After Shannon*: Proc. of the IEEE **88**, 569-587 (2000).
 - [17] B.B. Mandelbrot and J.W. Van Ness, *Fractal Brownian motions, fractional noises and applications*: SIAM Rev. **10**, 422-437 (1968).
 - [18] *The Science of Fractal Images*, edited by H.O. Peitgen and D. Saupe (Springer-Verlag, New York, 1987).
 - [19] Z.S. She, E. Jackson, and S.A. Orszag, *Structure and Dynamics of Homogeneous Turbulence: Models and Simulations*: Proc. Roy. Soc. London A **434**, Issue 1890, 101-124 (1991).
 - [20] B.L. Shraiman and E.D. Siggia, *Scalar turbulence*: Nature **405**, 639-646 (2000).
 - [21] J.A. Krommes, *Fundamental statistical descriptions of plasma turbulence in magnetic fields*: Phys. Rep. **360**, 1-352 (2002).
 - [22] A.N. Kolmogorov, *The Local Structure of Turbulence in Incompressible Viscous Fluid for Very Large Reynolds Numbers*: C.R. Acad. Sci. USSR **30**, 301-305 (1941).
 - [23] A. Corral, and M. Paczuski, *Avalanche Merging and Continuous Flow in a Sandpile Model*: Phys. Rev. Lett. **83**, 572-575 (1999).
 - [24] D.E. Newman, R. Sánchez, B.A. Carreras, and W. Farenbaugh, *Transition in the Dynamics of a Diffusive Running Sandpile*: Phys. Rev. Lett. **88**, 204304-1 (2002).